

XII. CALASTRAPIA, gen. nov.

Range. N.W. New Guinea.

Type.
C. splendidissima.

XIII. PARADISEA.

Paradisea, Linn. Syst. Nat. i. p. 166 (1766)

Type.
P. apoda.

Range. New Guinea, Salawati, Jobi, Misori, Mysol, Waigiou, Ghemien, Batanta, Aru Islands.

Key to the Species.

- a. Yellow of head confined to the crown and nape ; the back maroon from the nape to the tail.
- a'. Ornamental flank-plumes yellow ; no yellow collar below the green throat ; breast and abdomen not contrasting strongly with the velvety brown of the chest, which is only a little darker { 1. *apoda.*
2. *novæ-guineæ.*
- b'. Ornamental flank-plumes crimson ; a distinct yellow collar below the green throat ; a distinct patch of velvety brown on the chest, in contrast to the vinaceous chestnut of the breast and abdomen 3. *raggiana.*
- b. Yellow of head continued over the mantle and back.
- c'. Yellow extending over the whole back to the upper tail-coverts.
- a''. Green of forehead extending to the line of the eyes, and forming a broad frontal band ; a narrow yellow collar below the green throat ; underparts of the same tint, the chest not velvety in contrast to the breast ; flank-feathers yellow, washed with reddish-brown, white at the ends 4. *maricæ.*
- b''. Green of forehead confined to a narrow basal line ; a yellow collar below the green throat.
- a³. Chest velvety black, in contrast to the light vinous-brown breast.
- a⁴. Flank-feathers golden orange 5. *augustæ-victoriæ.*
- b⁴. Flank-feathers crimson 6. *intermedia.*
- b³. Chest delicate lilac-grey like the breast ; flank-plumes crimson, with a patch of silky black plumes near the base on each side of the abdomen 7. *decora.*
- d'. Yellow of back restricted to the interscapular region, the lower back and rump chestnut ; flank-plumes yellow 8. *minor* and allies.

XIV. TRICHOPARADISEA.

Trichoparadisea, Meyer, Abhandl. k. zool. Mus. Dresden, 1892-93, p. 20 (1893)

Type.
T. gulielmi.

Range. German New Guinea.

XV. URANORNIS.

Uranornis, Salvad. Ann. Mus. Civic. Genov. ix. p. 191 (1876)

Type.
U. rubra.

Range. Islands of Waigiou, Ghemien, and Batanta.

XVI. PARADISORNIS.

Paradisornis, Meyer, Zeitschr. ges. Orn. ii. p. 385 (1885)

Type.
P. rudolphi.

Range. S.E. New Guinea.

XVII. LAMPROTHORAX.

Lamprothorax, Meyer, Abhandl. k. zool. Mus. Dresden, 1894-95, no. 2, p. 3 (1896)

Type.
L. wilhelminæ.

Range. N.W. New Guinea.