

1. *paradisea*.  
*Drepananax*, Sharpe, Bull. Brit. Orn. Club, iv. p. xv (1894) . . . . .  
*Range*. New Guinea.  
 Type. *D. bruijni*.

2. *victoriae*.  
*Epimachus*, Cuvier, Règne Anim. i. p. 407 (1817) . . . . .  
*Cinnamolegus*, Less. Hist. Nat. Ois. Parad. Syn. p. 32 (1833) . . . . .  
*Range*. New Guinea.  
 Type. *E. speciosus*.

1. *magnifica*.  
*C. magnifica*.  
*a*. Tail with a steel-blue or steel-green gloss ; back spangled with metallic-green or steel-blue.  
*a'*. Elongated flank-plumes black ; under surface of body black, with an oily-olive gloss, slightly inclining to purplish . . . . .  
*b'*. Elongated flank-plumes brown ; under surface of body brown, with a distinct olive shade or purplish gloss . . . . .  
*c'*. Elongated flank-plumes green, like the abdomen, contrasting with the emerald-green chest ; throat black, followed by a coppery-red thorax . . . . .  
*b*. Tail and wings with a violet-purple gloss ; spangles on back reddish-purple . . . . .  
 Type. *E. speciosus*.  
 1. *speciosus*.  
 2. *meyeri*.  
 3. *astrapioides*.  
 4. *ellioti*.

## SUB-FAMILY II. PARADISEINÆ.

## Key to the Genera.

- a*. No fleshy-yellow wattle round the eye.  
*a'*. Tail of extreme length and graduated, the central feathers broad like the others, exceeding the length of the body of the bird, and not reduced to narrow plumes or wire-like shafts.  
*a''*. Tail twice the length of the body ; round the hind-neck a metallic frill . . . . .  
*b''*. Tail four times the length of the body of the bird ; no metallic frill round the hind-neck . . . . .  
*c''*. Tail shorter, scarcely exceeding the length of the body of the bird ; no metallic frill round the hind-neck . . . . .  
*b*. Tail square or slightly rounded, with the two central feathers enormously elongated and reduced to shaft-like plumes, either very narrow and slightly spatulate at the ends (*Paradisornis*) or like wire (*Paradisea*) or whalebone (*Uranornis*), with the webs of the long flank-feathers disconnected near their tips and without radii.  
*d''*. Flank-plumes very long, full, and ornamental, reaching beyond the tail by more than the wing's length . . . . .  
*e'*. Flank-plumes ornamental, but not reaching beyond the tail by as much as the wing's length.  
*a<sup>3</sup>*. Centre tail-feathers as in *Paradisea*, but the flank-plumes not so dense and having thread-like ends like a spider's web ; on the throat a metallic-green shield . . . . .  
*b<sup>3</sup>*. Centre tail-feathers reduced to horny shafts like whalebone, with no barbs or web of any kind ; flank-plumes red, with thread-like ends ; a metallic-green shield on the throat . . . . .  
 Type. *H. mantouii*.  
 Type. *P. duivenbodii*.  
 Type. *I. bensbachii*.  
 Type. *S. nigricans*.  
 Type. *S. nigricans*.  
 Type. *D. albertisi*.  
 Type. *D. albertisi*.  
 1. *albertisi*.  
 2. *geisleri*.  
 3. *cervinicauda*.  
 10. *ASTRAPIA*.  
 11. *ASTRARCHIA*.  
 12. *CALASTRAPIA*.  
 13. *PARADISEA*.  
 14. *TRICHOPARADISEA*.  
 15. *URANORNIS*.