

AMADINA JAHUANA

General History

The first specimens were taken by Gmelin, at the mouth of the Amazon, in 1769. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers.

The first specimens were taken by Gmelin, at the mouth of the Amazon, in 1769. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers.

The first specimens were taken by Gmelin, at the mouth of the Amazon, in 1769. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers.

The first specimens were taken by Gmelin, at the mouth of the Amazon, in 1769. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers.

The first specimens were taken by Gmelin, at the mouth of the Amazon, in 1769. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers.

The first specimens were taken by Gmelin, at the mouth of the Amazon, in 1769. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers. It is a very common bird in the mountains of the Andes, where it is found in the most fertile and cultivated parts of the country. It is also very common in the plains and on the banks of the rivers.