

PETROICA GOODENOVII, *Jard. and Selb.*

Red-capped Robin.

Muscicapa Goodenovii, Vig. and Horsf. in Linn. Trans., vol. xv. p. 245.—Jard. and Selb. Ill. Orn., vol. i. pl. 8, fig. 2.

Petroica Goodenovii, Jard. and Selb. Ill. Orn., Add., vol. ii.—Gould, Syn. Birds of Australia, Part I.

Mé-ne-ge-dang, Aborigines of the mountain districts of Western Australia.

Red-capped Robin of the Colonists.

THE red crown and much smaller size of this beautiful Robin at once distinguishes it from every other species of the genus yet discovered. Although not plentiful in any part I have visited, it is very generally distributed over the whole of the southern portion of Australia. I have observed it myself in South Australia and in New South Wales, and Mr. Gilbert killed it in Western Australia, where, however, it is very local, for he only met with it in two spots, one in the York district and the other at Kojenup, about one hundred miles towards the interior from King George's Sound. I have not yet heard of its being an inhabitant of the northern portion of the country.

I generally observed it either singly or in pairs, and it appeared to give a decided preference to the beds of dry rivulets, and to thinly timbered plains, the dense brushes near the coast never being visited by it; it would seem therefore to be a species peculiar to the interior of the country.

The whole of the actions and economy of this bird so closely assimilate to those of the *Petroica multicolor*, that it is unnecessary to repeat a description of them here; of its nidification no information has yet been obtained; but in this respect also it doubtless closely resembles the same species.

It possesses a peculiarly sweet and plaintive song, very much like that of the European Robin, but more weak and not so continuous.

Its food consists of insects of various kinds.

The male has the upper surface, neck, upper part of the breast and wings brownish black; wing-coverts and secondaries edged with white, forming a broad stripe along the wings; middle of the outer web of the quills with a narrow white margin; forehead, crown, and lower part of the breast bright scarlet, passing into white on the vent; irides, bill and feet blackish brown; soles of the feet yellow.

The female, as is the case with the females of the other species, differs much from her mate in the colouring of the plumage, which difference will be more clearly perceived in the accompanying illustration than by the most minute description.

The Plate represents the two sexes of the natural size.