

Subfamily ALAUDINÆ, Bonap.

Genus MIRAFRA, Horsf.

One, if not two, species of this well-defined genus inhabit Australia. At present one only has been characterized; but the bird of this form, frequenting the intertropical portions of the country, may prove to be a distinct species.

230. Mirafra Horsfieldii, Gould Vol. III. Pl. 77.

Family FRINGILLIDÆ, Vig.

The Finches of Australia comprise twenty well-marked species, referable to several genera or subgenera, each of which exhibit a slight difference in structure, accompanied, as is always the case, by a difference in habit, and in the districts inhabited; thus the true *Estrelidæ* frequent grassy patches in the glades of the forests, the open parts of gullies, &c.; the *Amadinae*, the stony hills and flats; the *Poëphila*, the grass beds of the open plains; and the *Donacolæ*, the grasses of the marshy districts and reed-beds: of the habits of *Emblema* nothing is known; its pointed bill indicates some peculiarity in its economy differing from those of the other genera.

All the species build, I believe, large grassy nests with a spout-like opening.

The whole of those figured are peculiar to Australia.

Genus ESTRELEDA, Swains.

- | | |
|--|-------------------|
| 231. Estrelida bella | Vol. III. Pl. 78. |
| <i>Loxia nitida</i> , Lath. Gen. Hist. vol. v. p. 268? | |
| 232. Estrelida oculea | Vol. III. Pl. 79. |
| 233. Estrelida Bichenovii | Vol. III. Pl. 80. |
| 234. Estrelida annulosa, Gould | Vol. III. Pl. 81. |
| 235. Estrelida temporalis | Vol. III. Pl. 82. |
| 236. Estrelida Phaëton, Gould | Vol. III. Pl. 83. |
| 237. Estrelida ruficauda, Gould | Vol. III. Pl. 84. |
| 238. Estrelida modesta. | |
| <i>Amadina modesta</i> , Gould | Vol. III. Pl. 85. |

Genus AMADINA, Swains.

- | | |
|--|-------------------|
| 239. Amadina Lathami | Vol. III. Pl. 86. |
| 240. Amadina castanotis, Gould | Vol. III. Pl. 87. |

Genus POËPHILA, Gould.

Generic characters.

Bill considerably swollen at the base, rendering it nearly as deep and broad as it is long; *wings* moderately long, the first quill rudimentary, the four next equal in length; *feet* plantigrade, *toes* slender; the middle toe much longer than the lateral ones, which are equal in length; *hind-toe* much shorter than the middle one; *tail* strictly cuneiform, the two middle feathers much produced.

- | | |
|---|-------------------|
| 241. Poëphila Gouldiæ. | Vol. III. Pl. 88. |
| <i>Amadina Gouldiæ</i> , Gould | |
| 242. Poëphila mirabilis, Homb. et Jacq. | Vol. III. Pl. 89. |