MONTIFRINGILLA HÆMATOPYGIA, Gould.

Red-rumped Mountain Finch.

Montifringilla hæmatopygia, Gould in Proc. of Zool. Soc., Mar. 25, 1851.

The specimens from which the accompanying figures were taken were killed in Thibet by Lord Gifford; they belong to a species intimately allied to the *Fringilla* (*Linaria*) *Gebleri* of M. Brandt, but which differs from it in its larger size, in the darker colour of the head and face, and in the darker colouring of the back and rump, which latter is moreover ornamented with a patch of blood-red; it is also destitute of the orange-red mark on the shoulders, which M. Brandt states to be characteristic of his species.

I am indebted to Lord Gifford for the loan of this fine bird, which is second in interest only to the new and beautiful *Syrrhaptes Thibetanus*, also brought by his lordship, and of which figures will be found in their proper place in this work.

Face and forehead brownish black, gradually blending into the light greyish brown of the upper surface; rump stained with blood-red; upper tail-coverts brown, tipped with dull white; tail dark brown, each feather margined externally with white; wing-coverts hoary; wings dark brown, the first four primaries narrowly edged with white, the next five primaries with a broad streak of white along the basal portion of their external webs terminating in a line with the extremities of the secondaries, which are externally fringed with hoary; spurious wing dark brown, margined at the base with whitish; under surface very light brown, gradually becoming paler, until on the under tail-coverts the hue is buffy white; bill and feet bluish black.

CANDA WELL BUILDING TO THE COURT OF THE COUR

The Plate represents the two sexes of the natural size.