

HARPACTES REINWARDTI.

Reinwardt's Trogon.

SPECIFIC CHARACTER.

Mas. *Capite, dorso tectricibusque caudæ superioribus saturatè viridibus; pectore olivaceo-brunneo; ventre flavo, ad latera aurantiaco.*

Male.—Crown of the head, back, wing-, and upper tail-coverts dark shining green; centre of the wing dark green, crossed by numerous narrow wavy lines of dull yellow; primaries greyish black, margined externally with white; six middle tail-feathers dark glossy green; the three lateral feathers on each side of the same hue at the base of the outer web, and greenish slate-colour on the inner, the outer feather margined with white throughout the entire length of its outer web, the next on each side from near the base to the tip, and the third for about half its length from the tip, all three tipped with white; throat pale orange-yellow; across the breast a broad band of yellowish olive-brown; breast and all the under surface rich orange-yellow; bill deep red; orbits cobalt-blue; space between the orbits and bill pea-green; feet orange.

Total length from $12\frac{1}{2}$ to $13\frac{1}{2}$ inches, *bill* $\frac{1}{2}$, *tail* $7\frac{1}{2}$, *wing* $5\frac{1}{2}$.

Female.—Similar in colour; but the green of the head is duller; the throat is pale yellowish white; the band across the breast is greyish brown, without any of the yellow suffusion seen in the male; the orange of the abdomen is much paler; and the wavy lines of yellow on the centre of the wing are broader, paler, and more conspicuous.

Trogon Reinwardti, Temm. Pl. Col. 124.—Id. Mon. of Trog. in Pl. Col.—Less. Traité d'Orn., p. 122.—Gould, Mon. of Trog., pl. 27.

——— *Reinwardtii*, Steph. Cont. of Shaw's Gen. Zool., vol. xiv. part 1, p. 221.

Apaloderma Reinwardtii, Swains. Class. of Birds, vol. ii. p. 337.

——— *Reinwardtii*, Gould, Mon. of Trog., syn. spec. *Apaloderma*, sp. 2.

Trogon (Apaloderma) Reinwardtii, Gould, Mon. of Trog., List of Plates, sp. 27.

Harpactes Reinwardtii, Hartl. Verz. Mus. Brem., p. 12.—Gray and Mitch. Gen. of Birds, vol. i. p. 71, *Harpactes*, sp. 9.—Gray, List of Spec. of Birds in Coll. Brit. Mus., pt. ii. sect. i. *Fissirostres*, p. 45.—Bonap. Consp. Gen. Av., tom. i. p. 151, *Harpactes*, sp. 9.

——— *Reinwardtii*, Gould, Birds of Asia, part xvii. pl. 4.

Apalharpactes Reinwardtii, Bonap. Consp. Vol. Zygod. p. 14. gen. 5 e. 43.

"*Trogon sulphureus*, Begbie," Bonap. Consp. Gen. Av. tom. i. p. 151, in syn. *Harpactes*, sp. 9.

Hapalarpactes Reinwardti, Cab. et Heine, Mus. Hein., Theil iv. p. 162.

Trogon Reinwardti, Huxl. in Proc. Zool. Soc. 1867, p. 444.

THE native country of this fine bird is Java, where it was discovered by Professor Reinwardt; and Mr. Wallace informs me it inhabits the western parts of the country, up to an altitude of 6000 feet above the sea.