

SYNOPSIS OF THE TROGONIDÆ.

SPECIES OF THE NEW WORLD.

THE ORBITS ALWAYS COMPLETELY FEATHERED.

A. Bill moderately stout; maxilla with a single terminal notch; wing- and tail-coverts more or less elongated.

PHAROMACRUS.

*a*¹. Occipital crest well developed; four central tail-coverts far exceeding the rectrices. 1. *mocinno*.

*b*¹. Occipital crest moderate; tail-coverts shorter than, or hardly exceeding the rectrices.

*a*². Three outer rectrices terminally white.

*a*³. Occipital crest green; base only of outer rectrix black 2. *antisianus*.

*b*³. Occipital crest golden green; basal half of outer rectrix black 3. *fulgidus*.

*b*². Rectrices black.

*c*³. Bill yellow; coverts a little longer than rectrices 4. *auriceps*.

*d*³. Bill red; coverts shorter than rectrices 5. *pavoninus*.

B. Bill feeble; maxilla with a single terminal notch; wing-coverts not elongated; tail-coverts only slightly so; head with elongated lateral plumes EUPTILOTTIS.

A single species (*E. neoæneus*).

C. Bill moderate; edges of both mandibles strongly serrated; wing- and tail-coverts not elongated; tail-feathers without prolonged terminal emarginations; wing-coverts either wholly black or minutely freckled with black and grey; throat and chest either green or blue or greyish black; no subterminal black spot on outer web of the two outer rectrices on each side TROGON.