

AND

Sa

And. fronte, genis, gulâ, corpore

Front and sides of the head, ch
wings, tail, and thighs yellow
mediately in front of the orbits
the middle of the bill; apical halves
blue; their apical halves
orbita red; irides lemon-yellow.

Total length, 15½ inches; bill

L'Araçari Baillon, Levaill. I

Ramphastos Bailloni, Vieill.

Méth. Orn., part

Pteroglossus Bailloni, Wagl.

—Gray and Mi

Consp. Gen. Av.

glossus, sp. 7.

—*croceus, Jard. & S*



Most of the travellers who have s
liancy of the colouring of their soft
parts of their bodies, some being r
ornithologist of distant countries c
of the gay and brilliant hues which
trouble, that I have been enabled to
correctly. In no instance do these
in fact is the difference of appear
were we not aware of the changes
identical. The intensity of the col
at others; this may be due to the
brilliant during the pairing and bre
this species I am indebted to the
London on his return to Vienna, a
with drawings and full particulars

As this bird is not characterized
which are to be found in all the true
of a uniform tint like the members
its most natural situation to be in
Brazil, over which it is rather wide
woods near Antonio Dias, on the S
lives in pairs, and that the stomach

Either the sexes differ very con
varieties; some individuals havin
swollen. Could I detect any differe
regard them as two species, but are
sufficient to call the attention of
bird, to the subject.

The Plate represents the bird o

ANDIGENA BAILLONI.

Gould & Richter del. & lith.
Ballard & Watson, Imp.

