LYNCORNIS TEMMINCKII, Gould.

Lync. vertice cinerescenti-fusco, nigro exilitèr fasciato, et apud plumarum apicem maculato; gulà castaneo nigroque fasciatà; annulo collum circumdante, anticè albo, posticè cervino; capitis penicillis, brevibus, nigris; pectore nigrescente, castaneo maculato; dorso, castaneo, cervino, maculisque nigris, ornato; remigibus alarum nigrescenti-fuscis, castaneo obsoletè fasciatis; abdomine pallidè cervino, plumis singulis ad basin nigris; caudà fuscà, castaneo interruptè fasciatà; rostro pedibusque fuscis.

Long. tot. $10\frac{1}{2}$ unc.; rostri, $1\frac{1}{8}$; alæ, 8; caudæ, $5\frac{1}{2}$; tarsi, $\frac{1}{2}$.

Crown of the head greyish brown, minutely barred with black, many of the feathers having a black spot at the apex; throat barred with rich chestnut and black; round the neck a collar, which is white anteriorly, and buff posteriorly; egrets short and black; chest blackish, freckled with chestnut; back ornamented with tints of chestnut and fawn, spotted and irregularly freckled with black; quills blackish brown, with nearly obsolete bars of chestnut; under-surface pale buff mingled with black, which latter colour occupies the base of each feather; tail brownish black, with interrupted bars of chestnut; bill and feet brown.

Lyncornis Temminckii, Gould, in Proc. of Zool. Soc. Part VI. 1838.

This species I had named *Lyncornis collaris* in my MS., and my friend M. Temminck has recently sent me a specimen under the name of *imberbis*; now as the term *collaris* is applicable to every species of the genus, and the term *imberbis* equally so, I have considered it best to rescind both these specific appellations, and to substitute in their place that of *Temminckii*, thus dedicating the species to my excellent friend.

This appears to be a much more abundant species than the two preceding; I have several examples in my own collection. It is a native of Borneo.

The figure is of the natural size.