

## Genus AMBLYPTERUS, Gould.

### CHARACTERES GENERICI.

*Rostrum debile et elongatum. Nares elevatae et rotundatae. Rictus setis robustis instructus, rostro longioribus. Alae truncatae; remigibus externis sextis ferè aequalibus et falcatis; remigibus secundo, tertio, quarto ad externum pogonium emarginatis, septimo, octavo, nono ad apices elongatis et attenuatis, decimo abruptè brevi; secondariis brevissimis, rotundatis, tertiarisque elongatis, conditis. Cauda brevissima et quadrata. Pedes ambulatorii. Tarsi elongati, graciles, squamis indistinctis anticè et posticè fasciati; digito intermedio longissimo et gracillimo; digitis lateralibus brevibus et aequalibus; digito postico parvo, debili et libero; unguibus elongatis, ungue medio pectinato.*

## AMBLYPTERUS ANOMALUS, Gould.

*Amb. summo capite, corpore suprà, et alis cinereo-fuscis, singulis plumis nigro irregulariter sparsis et maculatis; primariis nigris, ad bases rubrescenti-cervinis, ad apices albis; secondariis cervinis, nigrescenti-fusco irregulariter fasciatis; rectricibus caudæ cervinis, nigrescenti-fusco irregulariter fasciatis et maculatis; duabus centralibus cinereo-fuscis; gutture, pectore, et abdomine ad partem superiorem nigrescenti-fuscis, singulis plumis cervino maculatis; abdomine imo pallidè cervino, singulis plumis nigrescenti-fusco transversim fasciatis; rostro fusco; pedibus pallidè fuscis.*

Long. tot.  $6\frac{3}{4}$  unc.; rostri, 1; alæ,  $5\frac{2}{3}$ ; caudæ, 3; tarsi,  $\frac{7}{8}$ .

A pale fawn-coloured stripe from the nostrils over each eye; crown of the head, back, and wings greyish brown; the centre of each feather irregularly dashed and spotted with black; primaries black, with the basal portion reddish fawn colour, and the tips white; secondaries fawn colour, irregularly barred with blackish brown, the tips being paler; two centre tail-feathers greyish brown, irregularly barred and speckled with blackish brown; the remainder of the feathers fawn colour, irregularly barred with blackish brown, and speckled on the intermediate spaces with the same colour, the whole largely tipped with light fawn; throat, chest, and upper part of the abdomen blackish brown, each feather on the chest having a longitudinal stripe of fawn down the centre, the whole minutely speckled with fawn; lower part of the abdomen light fawn colour, each feather transversely barred with blackish brown; bill brown; feet light brown.

*Amblypterus anomalus*, Gould, in Proc. of Zool. Soc. Part V. 1837, p. 105.

---

A SINGLE specimen of this truly anomalous Goat-sucker forms part of the collection of the British Museum; it is the only one I have ever seen, and is, I believe, entirely new to science.

The only information I have been able to obtain respecting it is, that Mr. J. E. Gray purchased the specimen above mentioned from a collection of skins at Liverpool, which were said to be from Demerara.

The figure is of the natural size.