

REPRODUCTION OF THE

Long-billed Curlew

REPRODUCTION OF THE LONG-BILLED CURLEW. The species referred to in this paper is the Long-billed Curlew, *Numenius longirostris*, Linn. It is a large wading bird, with long legs and a long, straight bill. It is found in marshy and coastal areas, and is common in the United States and Canada. The Long-billed Curlew is a migratory bird, and it breeds in the northern part of its range. It is a monogamous species, and it builds a nest of mud and sticks. The female lays a clutch of four or five eggs, and she incubates them for about three weeks. The young are born naked and helpless, and they stay in the nest for about three weeks before they leave. The Long-billed Curlew is a game bird, and it is hunted for food and sport. It is also a pest to farmers, because it eats the crops. The Long-billed Curlew is a beautiful bird, and it is a pleasure to watch it in its natural habitat.

It is the purpose of this paper to describe the reproduction of the Long-billed Curlew. The species referred to in this paper is the Long-billed Curlew, *Numenius longirostris*, Linn. It is a large wading bird, with long legs and a long, straight bill. It is found in marshy and coastal areas, and is common in the United States and Canada. The Long-billed Curlew is a migratory bird, and it breeds in the northern part of its range. It is a monogamous species, and it builds a nest of mud and sticks. The female lays a clutch of four or five eggs, and she incubates them for about three weeks. The young are born naked and helpless, and they stay in the nest for about three weeks before they leave. The Long-billed Curlew is a game bird, and it is hunted for food and sport. It is also a pest to farmers, because it eats the crops. The Long-billed Curlew is a beautiful bird, and it is a pleasure to watch it in its natural habitat.