

Thalurania furcata, Reichenb. Aufz. der Col. p. 7.
Coeligena furcata, Reichenb. Troch. Enum. p. 3, pl. 682. figs. 4498-99.
Trochilus furcatus, Burm. Th. Bras. tom. ii. p. 335.
Habitat. Cayenne and Guiana.

117. THALURANIA FURCATOIDES, Gould.

Thalurania furcatoides, Gould, in text to *T. furcata*.
Habitat. Para and the lower part of the Amazon.

This bird is very like *T. furcata*, but is of smaller size, has a much less-forked tail, and the breast ultramarine-blue instead of purplish-blue. I have at this moment seven specimens before me, all of which are alike, and readily distinguishable from the Cayenne bird.

118. THALURANIA FORFICATA, Cab.

Thalurania forficata, Cab. et Hein. Mus. Hein. Theil iii. p. 24.
Habitat. Supposed to be the neighbourhood of Para.

Through the kindness of Dr. Peters, the Director of the Royal Museum of Berlin, I have had their specimen of *T. forficata* sent to me for comparison. It is certainly distinct from any of the species contained in the collections of this country: in size it is nearly the same as *T. furcata*; but its bill is rather shorter, its tail longer, more deeply forked, and of a purplish-black colour instead of steely-black: it differs from that bird also in having the blue colouring more extended down the back, approaching to that of *T. Watertoni*; the green of the throat is circumscribed and truncate below, as in *T. furcata*; and the crown of the head is black, but near the centre is a single small blue feather: I think it likely that this is accidental, as the bird appears to be fully adult: the under tail-coverts are black.

119. THALURANIA REFULGENS, Gould Vol. II. Pl. 102.
Habitat. Unknown; supposed to be Venezuela.

120. THALURANIA TSCHUDI, Gould Vol. II. Pl. 103.
Trochilus furcatus, Tschudi.
Habitat. Ucayali and the eastern part of Ecuador.

121. THALURANIA NIGROFASCIATA, Gould Vol. II. Pl. 104.
Habitat. Upper Amazon and Rio Napo.

122. THALURANIA VENUSTA, Gould Vol. II. Pl. 105.
Coeligena venusta, Reichenb. Troch. Enum. p. 3, pl. 683. figs. 4504-5.
 ——— *puella*, Reichenb. ib. p. 3.
Habitat. Chiriqui in Costa Rica.

123. THALURANIA COLUMBICA Vol. II. Pl. 106.
Thalurania columbica, Cab. et Hein. Mus. Hein. Theil iii. p. 24.
Habitat. The hilly parts of New Granada.

124. THALURANIA VERTICEPS, Gould Vol. II. Pl. 107.
Chlorestes verticeps, Reichenb. Troch. Enum. p. 4, pl. 705. fig. 4590.
Habitat. Ecuador.

125. THALURANIA FANNLE.

Coeligena Fanny, Reichenb. Troch. Enum. p. 3, pl. 683. figs. 4502-3.
Thalurania Fanny, Bonap. Rev. et Mag. de Zool. 1854, p. 254; Cab. et Hein. Mus. Hein. Theil iii. p. 23.
Habitat. The Andes of Quindios.

I have a specimen of this bird, collected by Warszewicz on the Cordillera of Quindios, which is precisely the same as the type specimen of the *T. Fannyi* of MM. Bourcier and Delattre. It differs from my *T. verticeps* in being considerably smaller and in having the abdomen purple-blue in lieu of cold prussian-blue.

126. THALURANIA ERIPHYLE Vol. II. Pl. 108.
Coeligena Eryphile, Reichenb. Troch. Enum. p. 3, pl. 582. figs. 4507-8.
Glaucoptis eriphyle, Burm. Th. Bras. tom. ii. p. 334.