of these grey and black-headed Wagtails, for which I must refer my readers to his valuable 'Iconografia

On reference to my MSS. I find the following note respecting this bird as observed by me in Norway, on reference to my MSS. I find the following note respecting this bird as observed by me in Norway, written while the matter was fresh in my memory. "From a careful examination of several specimens obtained on the Dovre-fjeld and in its neighbourhood, I am inclined to regard Budytes cinereocapilla and obtained on the Dovre-fjeld and in its neighbourhood, I am inclined to regard Budytes cinereocapilla and B. nigricapilla, Bonap. (M. melanocephala, Savi., nec Lichtenstein), as one and the same species. Males B. nigricapilla, Bonap. (M. melanocephala, Savi., nec Lichtenstein), as one and the same species. Males killed on approaching the Dovre-fjeld had dark grey heads, while those obtained near Jerkin had jet-black crowns. The dark-grey-headed birds had scarcely a trace of the stripe through the eye; the females, on the contrary, had the head olive-grey, a conspicuous stripe through the eye, and an indistinct mark of black on the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour from the grey to the black head is to be found on the Dovre-the chest. I believe every gradation of colour

As it is almost impossible to determine to which species the numerous notices that have appeared under the heading of Budytes cinereo-capilla have reference, I hesitate to quote any of them, with the exception of the following, from the pen of Lord Lilford, which, doubtless, pertains to the present bird. He states (in 'The Ibis' for 1860) that it "arrives in great numbers in Corfu about the middle of April, at which season it is to be found in small flocks in all the low meadows and maize-fields of the island. I never could find a nest of this species, though I have observed a few pairs during the whole summer."

In conclusion I may remark that the colouring and markings of the head of the male occasionally present slight differences, some specimens having the crown, lores, and ear-coverts black, others uniform dark sooty grey; and in others, again, the grey is broken by a faint white stripe above the ear-coverts, as seen in the Penzance example.

I must not omit to acknowledge the persevering kindness of Dr. Leith Adams, than whom no one of my friends rendered me more valuable assistance towards the illustration of these Wagtails. During his sojourn in Malta he collected for me a very large series, obtained during their passage from Africa to Europe in the spring, and on their return in autumn, at which latter season the young either accompanied the adults or followed them after a short interval. No examples of either the western Budytes Rayi or the eastern B. melanocephala were found among the birds I had the pleasure of receiving from Dr. Adams, the only species being B. cinereocapilla and B. flava.

The upper figure in the accompanying Plate represents the Penzance specimen; the two lower ones are from male and female examples shot by myself on the Dovre-fjeld, in Norway, on the 4th of July, 1856.