



OREOCINCLA AUREA.

J. Gould & H.C. Richter, del. et lith.

Walter, Imp.



Turdus aureus, Holla
 — *varius*, Pall.
 — *Whitei*, Eyt.
 — *varius* seu *WT*
Oreocincla Whitei, G.
Turdus immaculatus, Bl.
Oreocincla aurea, B.

Two known members
 the forests of most p
 in the extreme south
 and Japan; and it is
 Europe have probably
 The first recorder
 of British Birds, wh
 to give a figure of th
 Heron Court, near
 mentions that he wa
 who had allowed him
 of the forest-keepers
 it for his own collec
 to have been obtain
 had, about ten days
 of G. J. Allman, a
 chance for ornithol
 and sportsmen, Mr.
 an examination of w
 occurrence of White's
Turdus, and *Merula*
 "The opportunit
 rarely, that I thoug
 of any of its parts in
 Aon, where the s
 position is highly f
 and small enclosure
 the smaller birds—
 the autumn and win
 "In a small gra
 of mine observed a
 but soon recognize
 the account, I stim
 again flushed from
 again on the 26th
 it was busy in turn
 Blackbirds, Thrush
 and hawthorn berry
 disturbed, and ther
 one spot without
 feeding, was very u
 it happened to pass
 from these short
 Mr. Blyth says the
 situation the groun