## Family ANTHIDÆ.

## PIPITS.

RICHARD'S PIPIT.

An accidental visitant to the British Islands, where it is generally seen in winter and spring. The other countries frequented by it are Continental Europe, India, China, and in fact the whole of the temperate regions of the Old World.

Common in the champagne parts of France and other portions of Central Europe. Has been killed in England, and may therefore be regarded as forming part of our avifauna; but its visits must be considered purely accidental.

A stationary species, frequenting the shores and rocky parts of Britain—sparsely in Norfolk, but plentifully in Western Scotland and all the western islands, including the Outer Hebrides, Monach Isles, Haskar Rocks, and St. Kilda—keeping strictly to the sea-margin (R. Gray).

139. Anthus spinoletta . . . . . . . . . . . . . . . Vol. III. Pl. XI. Vinous Pipit.

A bird of France and the southern and eastern parts of Europe. Although it has been killed on our coast many times, it must be considered an uncertain and accidental visitor.

140. Anthus Ludovicianus.

See the remarks respecting this species in my account of the Vinous Pipit (Anthus spinoletta).

141. Anthus cervinus
Red-throated Pipit.

Vol. III, Pl. XII.

Mr. Harting has recorded, in the 'Field' for the 26th of August, 1871, the occurrence of two examples