

DOLCIN is NON-TOXIC . . . IT WILL NOT HURT THE HEART or other organs, even when taken in heavy dosage over protracted periods.

DOLCIN tablets, made under STRICTEST LABORATORY CONTROL, are TESTED for PURITY of ingredients and for Pharmaceutic uniformity.

DIRECTIONS

Read these directions CAREFULLY so that you may learn how you can get GREATEST BENEFIT from DOLCIN.

IT IS RECOMMENDED that not fewer than TWELVE DOLCIN tablets be taken daily (three tablets, with water, before each meal and three at bedtime) until acute symptoms are relieved. Then, follow this with EIGHT TABLETS (2 tablets, 4 times) daily for ten weeks . . . or until all symptoms disappear. At this time, cut the dosage to FOUR TABLETS (1 tablet, 4 times) for eight weeks more.

In Rheumatoid Arthritis, and in severe cases of any other type, best results have been had when the initial dosage is 24

tablets daily (6 tablets, 4 times) for four weeks, followed by 16 tablets a day for 4 weeks . . . then follow directions in preceding paragraph.

While it is generally better to take DOLCIN before meals, it makes no difference whether the DOLCIN at meal-time is taken before, during or after eating . . . BUT TAKE YOUR DOLCIN REGULARLY.

For most effective results it is very important to continue taking DOLCIN in reduced dosage for a few months after relief from pain has been obtained.

The DOLCIN formula is the only succinate-salicylate product with extensive scientific background of clinical and laboratory testing. Do not be misled by imitations which are claimed to be "just as good"!

Scientific Studies by outstanding authorities have established that the ingredients of DOLCIN provide effective relief of symptoms in the Rheumatic State. It is, therefore, suggested that DOLCIN be taken regularly, and for several months, after relief from pain has been obtained.